

NAME: Fonda Davis Eyler, Ph.D.
Professor Emerita, Department of Pediatrics
University of Florida College of Medicine
E-mail: eyler@peds.ufl.edu
2500 N.W. 23rd Terrace
Gainesville, FL 32605-2811

PUBLICATIONS (*Mentoring Author)

BOOKS

Contributor of Chapters

1. Eyler FD: Developmental assessment and care of the infant with respiratory problems. In: Eitzman DV, Koff PB, Kopotic R, Neu J (eds). Neonatal and Pediatric Respiratory Care. Annapolis, MD: C.V. Mosby, pp. 356-375, 1988.
2. Eyler FD: Developmental outcomes. In: Koff PB, Eitzman DV, Neu J (Eds). Neonatal and Pediatric Respiratory Care. St. Louis, MO: C.V. Mosby, pp. 440-501, 1993.
3. Woods NS, Behnke M, Eyler FD, Conlon M, Wobie K: Cocaine use among pregnant women: Socioeconomic, obstetrical, and psychological issues. In: Lewis M, Bendersky M (Eds). Mothers, Babies, and Cocaine: The Role of Toxins in Development. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc., pp. 305-332, 1995.
4. Eyler FD: Certain conditions originating in the perinatal period. In: Johnson SB, Perry NW, & Rozensky RH (Eds). Behavioral Applications to Medical Disorders. Volume 1 of the Handbook of Clinical Health Psychology, Boll TJ (ed). Washington, DC: American Psychological Association, pp. 483-531, 2002.

REFEREED PUBLICATIONS

Articles

1. Hofheimer JA, Poisson SS, Strauss ME, Eyler FD, Greenspan SI: Perinatal and behavioral characteristics of neonates born to multirisk families. J Dev Behav Pediatr 4:163-170, 1983
2. Resnick MB, Eyler FD, Nelson RM, Eitzman DV, Bucciarelli RL: Developmental intervention for low-birthweight infants: Improved early developmental outcome. Pediatrics 80:68-74, 1987
3. Behnke M, Carter RL, Hardt NS, Eyler FD, Cruz A, Resnick MB: The relationship of Apgar scores, gestational age, and birthweight to survival of low-birthweight infants. Am J Perinatol 4:121-124, 1987
4. Resnick MB, Reiss J, Eyler FD, Schauble P: Children's Developmental Services: A multidisciplinary program of psychological and educational services for neonatal intensive care. Journal of Counseling and Development 66:279-282, 1988

5. Behnke M, Eyler FD, Carter RL, Hardt NS, Cruz A, Resnick MB: Predictive value of Apgar scores for developmental outcome in premature infants. Am J Perinatol 6:18-21, 1989
6. Eyler FD, Courtway-Meyers C, Edens MJ, Hellrung DJ, Nelson RM, Eitzman DV, Resnick MB: Effects of developmental intervention on heart rate and transcutaneous oxygen levels in low birthweight infants. Neonatal Network 8:17-23, 1989
- *7. Adams C, Eyler FD, Behnke M: Nursing intervention with mothers who are substance abusers. J Perinatal Neonatal Nurs 3:43-52, 1990
8. Edens MJ, Eyler FD, Wagner JT, Eitzman DV: Neonatal ethics: Development of a consultative group. Pediatrics 86:944-949, 1990
9. Eyler FD, Delgado-Hachey M, Woods NS, Carter RL: Quantification of the Dubowitz Neurological Assessment of preterm neonates: Developmental outcome. Infant Behav Dev 14:451-469, 1991
- *10. Woods NS, Eyler FD, Behnke M, Conlon M: Cocaine use during pregnancy: Maternal depressive symptoms and infant neurobehavior over the first month. Infant Behav Dev 16:83-98, 1993
11. Behnke M, Eyler FD, Conlon M, Woods NS, Thomas V: The relationship between umbilical cord and infant blood gases and developmental outcome in very low birthweight infants. Clin Obstet Gynecol 36:73-81, 1993
12. Behnke M, Eyler FD: The consequences of prenatal substance abuse for the developing fetus, newborn, and young child. Internat J Addict 28:1341-1391, 1993
13. Lombardero N, Casanova OQ, Behnke M, Eyler FD, Bertholf RL: Measurement of cocaine and metabolites in urine, meconium, and diapers by gas chromatography/mass spectrometry. Ann Clin Lab Sci 23:385-394, 1993
14. Eyler FD, Behnke M, Conlon M, Woods NS, Frentzen B: Prenatal cocaine use: A comparison of neonates matched on maternal risk factors. Neurotoxicol Teratol 16:81-87, 1994
15. Behnke M, Eyler FD: Issues in prenatal cocaine use research: Problems in identifying users and choosing an appropriate comparison group. Inf Mental Health J 15:146-157, 1994
16. Casanova OQ, Lombardero N, Behnke M, Eyler FD, Conlon M, Bertholf RL: Detection of cocaine exposure in the neonate: Analyses of urine, meconium, and amniotic fluid from mothers and infants exposed to cocaine. Arch Pathol Lab Med 118:988-993, 1994
17. Behnke M, Eyler FD, Conlon M, Woods NS, Casanova OQ: Multiple risk factors do not identify cocaine use in rural obstetrical patients. Neurotoxicol Teratol 16:479-484, 1994
18. Eyler FD, Behnke M: Prenatal cocaine exposure: Consequences for child and family. J FL Med Assoc 82:603-606, 1995
19. Behnke M, Eyler FD, Conlon M, Casanova OQ, Woods NS: How fetal cocaine exposure increases neonatal hospital costs. Pediatrics 99:204-208, 1997

20. Winecker RE, Goldberger BA, Tebbett IR, Behnke M, Eyler FD, Wobie K, Karlix J, Conlon M, Bertholf R: Detection of cocaine and its metabolites in amniotic fluid and umbilical cord tissue. J Analyt Toxicol 21:97-104, 1997
21. Behnke M, Eyler FD, Woods NS, Wobie K, Conlon M: Rural pregnant cocaine users: An in-depth sociodemographic comparison. J Drug Issues 27:501-524, 1997
- *22. Wobie K, Eyler FD, Conlon M, Clarke L, Behnke M: Women and children in residential treatment: Outcomes for mothers and their infants. J Drug Issues 27:585-606, 1997
- *23. Wobie K, Eyler FD, Behnke M, Conlon M: Symbolic expression of feelings and depressive symptoms in high-risk mothers. J Fl Med Assoc 84:384-390, 1997
24. Eyler FD, Behnke M, Conlon M, Woods NS, Wobie K: Part I: Birth outcome from a prospective matched study of prenatal crack/cocaine use: Interactive and dose effects on health and growth. Pediatrics 101:229-237, 1998
25. Eyler FD, Behnke M, Conlon M, Woods NS, Wobie K: Part II: Birth outcome from a prospective matched study of prenatal crack/cocaine use: Interactive and dose effects on neurobehavioral assessment. Pediatrics 101:237-241, 1998
26. Behnke M, Eyler FD, Conlon M, Wobie K, Woods NS, Cumming W: Incidence and description of structural brain abnormalities in cocaine-exposed newborns. J Pediatr 132:291-294, 1998
- *27. Woods NS, Eyler FD, Conlon M, Behnke M, Wobie K: Pygmalion in the cradle: Observer bias against cocaine-exposed infants. J Dev Behav Pediatr 19:283-285, 1998.
28. Karlix JL, Behnke M, Eyler FD, Wobie K, Adams V, Freiburger B, Conlon M, Tebbett IR: Perinatal cocaine suppresses the fetal immune system. Pediatr Res 44:43-46, 1998
29. Behnke M, Eyler FD, Garvan CW, Tenholder M, Wobie K, Woods NS, Conlon M, Cumming W: Cranial ultrasound abnormalities identified at birth: Their relationship to perinatal risk and neurobehavioral outcome. Pediatrics 103:e4 1, 1-6, 1999
30. Eyler FD, Behnke M, Woods NS, Wobie K: Examiner masking in research on the effects of prenatal cocaine exposure. J Drug Issues 29:215-224, 1999
31. Eyler FD, Behnke M: Early outcome of infants exposed to drugs prenatally. Clinics in Perinatology 26:107-150, 1999
32. Behnke M, Eyler FD, Garvan CW, Wobie K: The search for congenital malformations in newborns with fetal cocaine exposure. Pediatrics 107:e74, 1-6, 2001
33. Winecker RE, Goldberger BA, Tebbett IR, Behnke M, Eyler FD, Karlix JL, Wobie K, Conlon M, Phillips D, Bertholf R: Detection of cocaine and its metabolites in breast milk. J Forensic Sci 46:1221-1223, 2001
34. Eyler FD, Behnke M, Garvan CW, Woods NS, Wobie K, Conlon M: Neonatal evaluations of toxicity and withdrawal related to prenatal cocaine exposure. Neurotoxicol Teratol 23:399-411, 2001

35. Behnke M, Eyler FD, Garvan CW, Wobie K, Hou W: Cocaine exposure and developmental outcome from birth to six months. Neurotoxicol Teratol 24: 283-295, 2002
36. Wobie K, Eyler FD, Garvan CW, Hou W, Behnke M: Prenatal cocaine exposure: An examination of mother-infant separation during the first year of life. J Drug Issues 34: 77-94, 2004
37. Hou W, Wang Z, Garvan CW, Behnke M, Eyler FD, Wu R: A generalized model for detecting genetic determinants underlying longitudinal traits with unequally spaced measurements and time-dependent correlated errors. Biostatistics 6: 420-433, 2005
38. Eyler FD, Behnke M, Garvan CW, Wobie K, Tebbett I: Relative ability of biologic specimens and interviews to detect prenatal cocaine use. Neurotoxicol Teratol 27: 677-687, 2005
39. Behnke M, Eyler FD, Warner TD, Garvan CW, Hou W, Wobie K: Outcome from a prospective, longitudinal study of prenatal cocaine use: Psychomotor development at three years of age. J Pediatr Psychol 31: 1-9, 2006
40. Behnke M, Eyler FD, Blackband SJ, Garvan CW, Leonard CM, Schmalfuss IM, Warner TD, Padgett KR: Neuroimaging in prenatally cocaine-exposed children. Neurotoxicol Teratol 28: 389-390, 2006
41. Warner TD, Behnke M, Eyler FD, Garvan CW, Hou W, Wobie K: Predicting behavior problems at three years in children with prenatal cocaine exposure: Structural equation modeling with a prospectively enrolled sample. J Behav Dev Pediatr 27: 83-92, 2006
42. Hou W, Garvan CW, Littell RC, Behnke M, Eyler FD, Wu R: A framework to monitor environment-induced major genes for developmental trajectories: Implications for a prenatal cocaine exposure study. Statistics in Medicine 25: 4020-4035, 2006
43. Warner TD, Behnke M, Eyler FD, Padgett KR, Hou W, Garvan CW, Leonard CM, Schmalfuss IM: Diffusion tensor imaging of frontal white matter and executive functioning in cocaine-exposed children. Pediatrics 118:2014-2024, 2006
44. Eyler FD, Warner TD, Behnke M, Hou W, Wobie K, Garvan CW: Executive functioning at ages 5 and 7 in children with prenatal cocaine exposure. Dev Neurosci 31:121-136, 2009
45. Warner TD, Behnke M, Eyler FD, Szabo N; Pre-adolescent cocaine use as determined by hair analysis in a prenatal cocaine exposure cohort. Neurotoxicol Teratol, 33(1): 88-99, 2011
46. Roussotte, F. Soderberg L. Warner, T. Narr, K. Lebel, C, Behnke, M, Eyler, FD, & Sowell, E. Adolescents with prenatal cocaine exposure show subtle alterations in striatal surface morphology and frontal cortical volumes. J Neurodev Disord. 4(33): 1-10 2013
[doi:10.1186/1866-1955-4-22](https://doi.org/10.1186/1866-1955-4-22) (PMID: 22958316)
47. Lebel, C., Warner, T., Colby, J., Soderberg, L., Roussotte, F., Behnke, M., Eyler, F. D., & Sowell, E. R. White matter microstructure abnormalities and executive function in adolescents with prenatal cocaine exposure. Psychiatry Research: Neuroimaging, 213(2): 161-168. 2013
doi.org/10.1016/j.pscychresns.2013.04.002 (PMID:23769420)

Invited Technical Reports

1. Eyler FD: Intervention with developmentally at-risk or delayed infants and their families: A guide for professionals. A training manual prepared for Developmental Services, HRS District III, State of Florida, January, 1-33, 1988.
2. Committee Member/Contributor: Toward a drug-free Florida: Report of the Drug-Exposed Infants/Families Committee, Governor's Drug Policy Task Force, Executive Office of the Governor, December, 1990.
3. Eyler FD, Maag L: The developmental outcome of children exposed prenatally to cocaine. Literature Review for Children's Medical Services Service Guidelines Panel.

Invited Articles

1. Cohen EA, Lidsky K, Eyler FD, Resnick MB: Considerations during intervention with premature infants. Newsletter, Am. O.T. Assoc., 6:1-2, 1983
2. Eyler FD: Effects of cocaine use on the fetus and mother. Intensive Caring Unlimited Newsletter 8(4): July/August 1990
3. Wobie K, Eyler FD, Behnke M, Conlon M, Woods NS: Symbolic expression of feelings and depressive symptoms in high-risk mothers. Florida Consortium of Newborn Intervention Programs Newsletter 6(3):7, 1993
4. Mehn M, Kazimour K, Eyler FD: A call to reinstate Florida's recently disenfranchised providers of early intervention services. Florida Consortium of Newborn Intervention Programs Newsletter 8(1):4-6, 1995
5. Behnke M, Eyler FD: Substance exposed infants. NAPSW Forum 24 (2): 8-11, 2004